

Title	Pharmacist-led academic detailing intervention in primary care: a mixed methods feasibility study
Authors	O'Riordan, David;Hurley, Eimir;Sinnott, Carol;Galvin, Rose;Dalton, Kieran;Kearney, Patricia M.;Halpin, James D.;Byrne, Stephen
Publication date	2019-01-22
Original Citation	O'Riordan, D., Hurley, E., Sinnott, C., Galvin, R., Dalton, K., Kearney, P. M., Halpin, J. D. and Byrne, S. (2019) 'Pharmacist-led academic detailing intervention in primary care: a mixed methods feasibility study', International Journal of Clinical Pharmacy. doi:10.1007/s11096-019-00787-6
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1007/s11096-019-00787-6
Rights	© 2019, Springer Nature Switzerland AG. This is a post-peer-review, pre-copyedit version of an article published in International Journal of Clinical Pharmacy. The final authenticated version is available online at: http://dx.doi.org/10.1007/s11096-019-00787-6
Download date	2023-05-05 01:47:47
Item downloaded from	http://hdl.handle.net/10468/7470



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Pharmacist-led academic detailing intervention in primary care: A mixed methods feasibility study

Introduction

The International Continence Society (ICS) has defined urinary incontinence as “*the complaint of any involuntary leakage of urine*” [1]. Despite the availability of evidence-based guidelines, some patients do not receive recommended treatment because of their reluctance to report this condition. In addition, when patients do seek help, many physicians are not familiar with the latest information on the appropriate methods of evaluating and treating patients with this condition [2].

Academic detailing is an interactive, convenient, and user-friendly approach that provides non-commercial evidence-based medical information tailored to the needs of an individual. A key tenant of the intervention is eliciting the provider's beliefs and attitudes and addressing those specifically during the encounter [3]. Academic detailers (who are usually pharmacists, nurses, or physicians) are trained to provide accurate, balanced, and up-to-date syntheses of the evidence on a clinical topic in an engaging format with healthcare professionals in their work environment [4]. These educational visits appear to be especially effective in improving prescribing appropriateness in general practice [5]. Other approaches to optimise prescribing behaviour include a healthcare professional such as a pharmacist carrying out medication reviews and providing feedback to GPs [6]. Computerised decision support systems (CDSS) are widely used tools that can support prescribing practice activities. They provide narrative information usually in the form of an alert at the time of prescribing or at the end of the patient consultation [7]. However, these approaches have their limitations, for example, there have been conflicting results reported for medication reviews conducted by pharmacists [8,9]. CDSS alerts can lead to alert fatigue, cognitive overload and desensitisation raising questions about the effectiveness of decision support [10]. This suggests that other interventions such as academic detailing are required to optimise clinical outcomes and patient care.

To date, no studies have evaluated the feasibility and acceptability of an academic detailing intervention with General Practitioners (GPs) in Ireland. Prior to implementing an intensive intervention like academic detailing on a national scale, it is prudent to evaluate the potential effect in a feasibility study. Eldridge *et al.* have defined a feasibility study as a study asking “*whether something can be done, should we proceed with it, and if so, how*”. They are used to estimate important parameters that are needed to design larger studies, for example: feasibility of recruitment, number of eligible participants, and selection of appropriate outcomes [11]. As the burden of urinary incontinence can impact on the lives of people worldwide (especially older adults), this would underline why a feasibility study of an academic detailing intervention is needed.

Aim of the study

The aim of this study was to assess the feasibility and acceptability to GPs of a pharmacist-led academic detailing intervention in Ireland.

Ethics approval

Ethical approval for this study was granted by the clinical research ethics committee of the Cork University Teaching Hospitals, Cork (reference ECM 4 (s) 10/05/16 & ECM 3 (dddd) 07/03/17) and the Mallow Primary Health Centre (MPHC), Cork Ethics Committee. Informed consent was obtained from all individual participants included in the study.

Methods

Study type

In this study a convergent parallel mixed methods design was used. This involves the separate collection and analysis of quantitative and qualitative data with the intent of merging the results of both analyses. The premise of a mixed methods approach is that the use of quantitative and qualitative methods in combination provides a better understanding of research problems than either approach alone [12]. The aim was to collect, analyse, and interpret integrated quantitative and qualitative data to assess the feasibility to GPs of an academic detailing intervention in primary care. The quantitative prescribing patterns of the GPs and their qualitative responses from the focus groups were integrated and synthesised.

Setting

This study was carried out in six General Practices in County Cork, Ireland. The primary researcher (D.O.R.) arranged a meeting with the lead GP in each practice, and a brief summary of the study was given. Other potential participants in each practice were contacted by telephone and invited to participate. Twenty-three GPs participated in the intervention. All GPs who participated in the study received a certificate of participation and a certificate for their continuing professional development (CPD).

Academic detailing training

D.O.R. received formal training in academic detailing by attending a two-day workshop at the National Resource Center for Academic Detailing (NaRCAD) in Boston, USA in May 2016. The workshop provided a critical foundation for the role as an academic detailer. It included sessions on the case for academic detailing and evidence-based medicine, planning a visit, use of educational materials and role plays.

71 **Academic detailing materials**

72 For the academic detailing sessions, materials developed by the Alosa Foundation were used.
73 The Alosa Foundation in Boston is a non-profit organization and is independent of the
74 pharmaceutical industry. It produces educational materials and decision making tools for
75 Academic Detailers, and provides training on academic detailing internationally. The Alosa
76 Foundation granted the authors permission to use their educational materials for this study. The
77 printed materials consist of evidence-based information on the prevalence of urinary
78 incontinence, an overview of the pharmacological and non-pharmacological interventions, cost
79 of drugs to treat the condition, and key messages.

80 **Choice of topic**

81 Prior to commencing GP recruitment, D.O.R. met with three GPs to discuss an intervention
82 topic. The topic of urinary incontinence was chosen for the academic detailing intervention by
83 GPs because they highlighted that it was a topic not discussed regularly among themselves,
84 and currently their only source of information is provided by pharmaceutical drug
85 representatives.

86 **Piloting the intervention**

87 The academic detailing intervention was piloted with three academic GPs in May 2016 before
88 the study was commenced. The GPs reported that they were very satisfied with this overall
89 educational approach. The topic delivered and the educational materials used during the visit
90 were also described as being very beneficial and relevant.

91 **Delivery of the intervention**

92 The intervention was delivered face-to-face with each GP in their practice. D.O.R contacted
93 the GP's receptionist in advance to book a 15-minute time slot with each GP for the meeting.

One of the tenets of academic detailing is to provide evidence based information over a succinct time period.

During the meeting, D.O.R provided the following key messages to GPs: (i) increase detection of and distinguish incontinence type to guide treatment, (ii) identify and rule out reversible causes of incontinence, (iii) encourage caffeine reduction, pelvic floor muscle training, and weight loss as first-line treatments, and (iv) judiciously prescribe medications for urgency symptoms (but not stress incontinence). GPs were encouraged to discuss these key messages with their patients when they presented for their next GP appointment. After the meeting, GPs were given a copy of the printed materials to use as a reference.

The academic detailing intervention was rolled out to participating GPs between June and September 2016. Supplementary Figure I shows the timeline of the study.

Quantitative method

A before and after analysis of patient medical records was conducted. All patients aged ≥ 65 years with urinary incontinence treated by participating GPs were included. These patients were identified by searching GP databases and notes. D.O.R. extracted data from medical records at five time points: six and three months before the intervention (T_{-6}), (T_{-3}), at the time of the intervention (T_0), and three and six months after the intervention (T_3), (T_6). For example, if the intervention was delivered to a GP in June 2016, data extraction and audit of their patient medical records was performed retrospectively in December 2016 for the following months: December 2015, March 2016, June 2016, September 2016 and December 2016. Patients were identified as having urinary incontinence based on a diagnosis by the GP or referral letters from consultant urologists. In some cases, prescription drugs (e.g. mirabegron, tolterodine) identified from medical records were used as proxies to indicate a diagnosis of urinary

incontinence. An assessment of patient medical records was carried out at five time points in the study to identify if pelvic floor muscle training was documented.

The following criteria were applied to the patient data recorded:

LUTS-FORTA criteria

Drugs to treat lower urinary tract symptoms (LUTS) in older people aged ≥ 65 years are classified on their appropriateness based on efficacy, safety, and tolerability using the Fit for The Aged (FORTA) criteria. These criteria classify drugs for the treatment of LUTS into four ordinal categories, A (absolutely: indispensable drug), B (beneficial: drugs with proven efficacy), C (careful: drugs with questionable efficacy/safety profiles), and D (don't: avoid in older people) [13].

The Drug Burden Index (DBI)

The DBI measures the cumulative exposure to anticholinergic and sedative medicines in older people and its impact on physical and cognitive function [14]. For each drug, the DBI ranges from 0-1, with 0 being no burden, 0.5 being exposure to the minimum daily dose, and upwards to 1 as the dose is increased exponentially [15]. In this study, the list of drugs with clinically significant anticholinergic and sedative effects were defined from a composite list developed from a review by Duran *et al.* 2013, the Anticholinergic Cognitive Burden (ACB) scale developed by Boustani *et al.* 2008, and from a study published by Ailabouni *et al.* 2017 [16-18]. This composite list consisted of 133 drugs (See Supplementary Table I).

Anticholinergic Cognitive Burden (ACB) scale

The cumulative effect of taking multiple medicines with anticholinergic properties is defined as the anticholinergic burden [19]. The ACB scale is based on a systematic literature review of medicines with known anticholinergic activity. The scale consists of 88 drugs with known anticholinergic activity and assesses individual drugs that have none, possible, or definite anticholinergic properties with a score ranging from 0 to 3 [17].

D.O.R. applied the LUTS-FORTA, DBI, and ACB to patient-related data retrieved from the electronic medical records. For validation purposes, the three types of criteria were applied independently by a second member of the research team to a random 10% sample of the data.

Outcomes

The three outcomes of interest were the overall changes in scores of LUTS-FORTA, DBI, and ACB in the patients.

Statistical analysis

It is important to note that a feasibility study is not a hypothesis testing study [20]. One of the key aspects of these studies is that they do not evaluate effectiveness as they are not powered to do so [21]. The main focus of this study was to assess feasibility of the intervention, and therefore the data were analysed using descriptive statistics. Continuous variables were presented as mean with standard deviation (SD) and range, or median with interquartile range (IQR), as appropriate, and categorical variables as frequency (percentage).

Qualitative method

After the intervention was delivered to participating GPs by D.O.R., focus groups were conducted with GPs to explore its feasibility and acceptability.

The focus groups were carried out by three researchers (E.H., C.S., and S.B.) between July and November 2016. A topic guide was developed based on discussion and consensus among all authors. The topic guide was iteratively refined after each focus group, was transcribed, and analysed to pursue emerging themes.

All focus groups were anonymised and fully transcribed and saved in QSR International NVivo Qualitative Data Analysis Software (V.10.22) to facilitate analysis. Data were analysed using thematic analysis. This flexible and useful research approach can potentially provide a rich and detailed account of the data [22].

Standardised reporting guidelines

The Good Reporting of a Mixed Methods Study (GRAMMS) framework was used to inform reporting of the findings [23]. (See Supplementary Table II).

Results

Quantitative results

The characteristics of 154 patients diagnosed with urinary incontinence included in the study are detailed in Table I. The documentation of pelvic floor muscle training was reported in 15% of patient medical records at only one-time point in the study (T₋₆).

LUTS-FORTA score

Figure I shows the number of patients prescribed drugs identified by the LUTS-FORTA criteria over time. According to the criteria, no patient was prescribed a drug rated in category A (absolutely: indispensable drug) at any time point. There was an increase in the number of patients prescribed drugs in category C (careful: drugs with questionable efficacy/safety profiles), over time, while there was no change in the number of patients prescribed drugs in category B (beneficial: drugs with proven efficacy) and D (don't: avoid in older people) over time. Supplementary Table III shows the drugs that were identified by the LUTS-FORTA criteria in this study.

Drug Burden Index

Almost 65% (100/154) of patients did not show any change in drug burden over time.

Anticholinergic cognitive burden (ACB) scale

Thirty-four percent of patients at T₋₆ months and 31% of patients at T₆ months had an ACB score of 0.

Qualitative results

Five focus groups were conducted in total (n=14 GPs). The mean number of participants per focus group was 3 (range 2 to 4). The focus groups ranged from 19 minutes to 48 minutes. The number of GPs working in a practice ranged from 1 to 7. The characteristics of GPs interviewed are detailed in Table II. Quotes supporting each theme/sub-theme are presented in Table III.

210 **Themes**

211 **Theme 1: The Academic detailing experience**

212 **Subtheme:** Convenience of academic detailing

213 Participants highlighted the convenience of the academic detailing session being carried out in
214 their working environment. They welcomed this educational visit being delivered with little
215 disturbance to their daily practice. They also reported that they are prepared to block out some
216 of their working time to accommodate this source of evidence-based information.

217

218 This was in contrast to the alternative sources of evidence-based information that are currently
219 available for GPs, for example: attending conferences, continuing medical education meetings,
220 or educational events on topics. Participants reported the frustration at not being able to attend
221 courses of interest due to the demands of their work schedule.

222

223 **Subtheme:** The interaction between participant and academic detailer

224 Participants described the interaction between the GP and the academic detailer as being
225 important to the success of the intervention. They reported that the session worked because it
226 felt relaxed and free of pressure. This was in contrast to their experience with some
227 pharmaceutical drug representatives who they described as having an aggressive approach
228 combined with an overload of information, which seemed to aggravate participants.

229

230

231

232 **Subtheme:** The educational materials

233 Participants said they liked the educational materials because they had a clear layout and were
234 easy to follow. They reported that they valued the succinct nature of the key messages, while
235 the tables and figures were presented in a straightforward way.

236

237 **Subtheme:** The topic: Urinary incontinence

238 The topic of urinary incontinence was agreed by a number of GPs prior to rolling out the study.
239 Participants reported that this topic was relevant and suitable to general practice. The relevance
240 of this topic facilitated the delivery of the intervention to GPs.

241

242 **Theme 2: Behaviour change**

243 Participants described the likelihood of changing their behaviour in treating patients with
244 urinary incontinence following the intervention. However, this change in behaviour could be
245 influenced by environmental resources, such as the availability of primary care
246 physiotherapists.

247

248 **Subtheme:** Knowledge gained

249 Participants were asked if they had gained any knowledge from the intervention. Some
250 participants of recent medical experience were not aware of the important role that non-
251 pharmacological methods play in treating urinary incontinence.

252

For some participants, the intervention served to refresh their knowledge with the topic rather than gain new knowledge as some of the treatment options may be more commonly used than others.

Theme 3: Sustainability

Subtheme: Academic detailing ownership

Participants were asked how this type of educational intervention could be rolled out to a wider group of GPs in Ireland. Some suggested that it could be affiliated with the Irish College of General Practitioners (ICGP), the professional and educational body for general practice in Ireland. The association with this recognised body could enhance the credibility of academic detailing among GPs.

Subtheme: Alternative formats of educational material

Participants suggested an online version of the educational material, which would be easier for them to manage in a setting where print materials over-accumulate or go missing.

Subtheme: Desire for practice staff involvement

Participants highlighted the importance of incorporating the wider members of the practice team in the academic detailing sessions. This is especially significant given the expanded role of nurses in primary care.

Subtheme: Future participation

Participants were asked if they would be interested in participating in future academic detailing studies. All indicated a willingness to do so.

Discussion

This study used a mixed methods approach to explore the feasibility and acceptability to GPs of a pharmacist-led academic detailing intervention.

Participants described the possibility of behaviour change following the intervention; however, this was partly dependent on the availability of primary care resources such as physiotherapists.

In Ireland, public patients are often on a waiting list to attend primary care physiotherapists.

Pelvic floor muscle training is an effective treatment for women with urinary incontinence and these exercises are often demonstrated by physiotherapists [24]. Therefore, if academic

detailing interventions are successful, then the treatment modalities recommended by academic detailers need to be resourced. Participants described the educational materials as being high

quality. These materials contain evidence-based information on the prevalence of urinary incontinence, an overview of the interventions and cost of drugs to treat the condition, and key

messages which may be very beneficial for GPs. However, if this information is not easily retrievable for GPs then they may not be used as a treatment resource during a patient

consultation [25]. This may have limited the change in prescribing outcomes in the study.

Participants called for these materials to become available as an online resource as they could be of value in optimising the diagnosis and management of urinary incontinence.

Comparison with existing literature

Allen *et al.* used a mixed methods study (a questionnaire and semi-structured telephone interviews) to explore family physician perceptions of academic detailing and identified several factors that encourage its use. They were: the relevance of the topic, the evidence-based approach adopted, and the educational material used [26]. These findings are similar to those in our study. The GPs selected the topic of urinary incontinence for the intervention and described it as relevant. They welcomed the idea of evidence-based information being presented to them in their practice and they also described the educational materials used as being of high quality. Soumerai recommends that educational materials should be brief and clearly presented [27]. For this intervention, all GPs received one visit from the detailer, however research has indicated that frequent reinforcement visits can optimise behaviour change [28]. In this study, GPs indicated a willingness to participate in future academic detailing studies. Hartung *et al.* and Anthierens *et al* also reported similar findings [4,25].

The measures of prescribing assessed: the LUTS-FORTA, DBI, and ACB showed minimal or no change in their scores following the intervention. As a feasibility study, this study was not conducted with a view to effecting change in prescribing outcomes, rather to see if the intervention was feasible and acceptable to GPs and whether this preliminary research was appropriate for successful implementation in subsequent larger studies. Our findings suggest GPs are open to this type of intervention, and recruitment and follow-up are possible. A larger study is required to more accurately examine the potential for meaningful change in clinical care.

The choice of topic for the academic detailing intervention was an important influence on GPs' readiness to participate in the study. It was possible that GPs chose urinary incontinence because of the difficulty in minimising the side-effects with associated treatments e.g.

anticholinergic effects and the limited availability of pharmacological options. The choice of urinary incontinence thus addressed GPs' learning needs but made evaluation of changes in prescribing difficult as many of the treatment options highlighted to GPs were non-pharmacological. Alternative topics for example, hypertension, atrial fibrillation, opioids in chronic pain may provide more scope to examine subsequent change in prescribing behaviours.

Strengths and limitations

To enhance the validity of the quantitative results, a sample of the data were independently reviewed by two healthcare professionals. The focus groups were arranged with GPs over a five-month period and this facilitated prolonged engagement with the data.

Although 23 GPs participated in the intervention, only 14 were available to attend the focus groups. All participating GPs were contacted in advance about the focus groups; however, some were away on the scheduled date while others who agreed to participate had to cancel at the last minute due to clinical emergencies or late clinics. One solution to this issue would be to organise them outside of practice hours e.g. at continuing professional development meetings.

The findings from this study may be beneficial to other researchers when developing their own study designs as they may enhance their approach or avoid similar pitfalls [29]. In future studies, a follow-up visit could be arranged with the GPs after four to six weeks to reinforce the key messages from the first visit and to identify if they have been successfully implementing any suggested changes. It would also give the academic detailer an opportunity to answer any additional questions that the GPs may have. Implementing academic detailing on a broader scale may benefit from, a "train the trainer" approach. Instructors could train their colleagues and this would help to build a pool of competent academic detailers. Finally,

randomized controlled trials that are methodologically robust and have large sample sizes should be considered [30].

Conclusion

This mixed methods study explored the feasibility and acceptability to GPs of a pharmacist-led academic detailing intervention. Overall, participants reported that this evidence-based approach was beneficial and welcomed further visits. The selection of a relevant topic appeared to be an important aspect of their positive response. The printed educational materials were reported as being well-presented and easy to follow, however an online version was preferred. Our findings provide a useful platform for the evaluation of academic detailing in primary care on a larger scale. Further research is needed in a larger population to determine the impact on patient outcomes and the cost-effectiveness of academic detailing.

Acknowledgements

The authors would like to acknowledge all the GPs who agreed to participate in this study. Additionally, gratitude is expressed to Alosa Health, who developed the academic detailing intervention “Evaluating and managing urinary incontinence”, and granted the authors permission to use their educational materials for this study. Alosa Health is a US non-profit which specialises in academic detailing. This body evaluates the evidence on clinical topics and synthesises the information into a 'user-friendly' format to be used in the interaction between the academic detailer and the clinicians. They provide information to improve clinical decision making and have no affiliation with any pharmaceutical company.

369 **Conflicts of interest**

370 There are no conflicts of interest to declare.

371

372 **Funding**

373 This research was funded by the Health Research Board SPHeRE/2013/1.

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

References

1. Abrams P, Cardozo L, Fall M, Griffiths D, Rosier P, Ulmsten U, van Kerrebroeck P, Victor A, Wein A (2002) The standardisation of terminology of lower urinary tract function: report from the Standardisation Sub-committee of the International Continence Society. *Neurourol Urodyn* 21 (2):167-178
2. Levy R, Muller N (2006) Urinary incontinence: economic burden and new choices in pharmaceutical treatment. *Advances in therapy* 23 (4):556-573
3. Avorn J, Soumerai SB (1983) Improving drug-therapy decisions through educational outreach. A randomized controlled trial of academically based "detailing". *N Engl J Med* 308 (24):1457-1463. doi:10.1056/nejm198306163082406
4. Hartung DM, Hamer A, Middleton L, Haxby D, Fagnan LJ (2012) A pilot study evaluating alternative approaches of academic detailing in rural family practice clinics. *BMC Fam Pract* 13:129. doi:10.1186/1471-2296-13-129
5. Grol R, Grimshaw J (2003) From best evidence to best practice: effective implementation of change in patients' care. *Lancet* 362 (9391):1225-1230. doi:10.1016/s0140-6736(03)14546-1
6. Bryant LJM, Coster G, Gamble GD, McCormick RN (2011) The General Practitioner-Pharmacist Collaboration (GPPC) study: a randomised controlled trial of clinical medication reviews in community pharmacy. *The International Journal of Pharmacy Practice* 19 (2):94-105. doi:10.1111/j.2042-7174.2010.00079.x
7. Pearson SA, Moxey A, Robertson J, Hains I, Williamson M, Reeve J, Newby D (2009) Do computerised clinical decision support systems for prescribing change practice? A systematic review of the literature (1990-2007). *BMC Health Services Research* 9:154. doi:10.1186/1472-6963-9-154
8. Kolhatkar A, Cheng L, Chan FK, Harrison M, Law MR (2016) The impact of medication reviews by community pharmacists. *Journal of the American Pharmacists Association : JAPhA* 56 (5):513-520.e511. doi:10.1016/j.japh.2016.05.002
9. Hanlon JT, Weinberger M, Samsa GP, Schmader KE, Uttech KM, Lewis IK, Cowper PA, Landsman PB, Cohen HJ, Feussner JR (1996) A randomized, controlled trial of a clinical pharmacist intervention to improve inappropriate prescribing in elderly outpatients with polypharmacy. *Am J Med* 100 (4):428-437. doi:10.1016/s0002-9343(97)89519-8
10. Ancker JS, Edwards A, Nosal S, Hauser D, Mauer E, Kaushal R, with the HI (2017) Effects of workload, work complexity, and repeated alerts on alert fatigue in a clinical decision support

426 system. BMC Medical Informatics and Decision Making 17 (1):36. doi:10.1186/s12911-017-
 427 0430-8

428 11. Arain M, Campbell MJ, Cooper CL, Lancaster GA (2010) What is a pilot or feasibility
 429 study? A review of current practice and editorial policy. BMC medical research methodology
 430 10 (1):67

431 12. Creswell JW, Plano Clark VL (2011) Designing and Conducting Mixed Methods Research.
 432 Sage
 433 [https://books.google.ie/books/about/Designing_and_Conducting_Mixed_Methods_R.html?id](https://books.google.ie/books/about/Designing_and_Conducting_Mixed_Methods_R.html?id=YcdlPWPJRBcC)
 434 [=YcdlPWPJRBcC](https://books.google.ie/books/about/Designing_and_Conducting_Mixed_Methods_R.html?id=YcdlPWPJRBcC)

435 13. Oelke M, Becher K, Castro-Diaz D, Chartier-Kastler E, Kirby M, Wagg A, Wehling M
 436 (2015) Appropriateness of oral drugs for long-term treatment of lower urinary tract symptoms
 437 in older persons: results of a systematic literature review and international consensus validation
 438 process (LUTS-FORTA 2014). Age Ageing 44 (5):745-755. doi:10.1093/ageing/afv077

439 14. Hilmer SN, Mager DE, Simonsick EM, Cao Y, Ling SM, Windham BG, Harris TB, Hanlon
 440 JT, Rubin SM, Shorr RI, Bauer DC, Abernethy DR (2007) A drug burden index to define the
 441 functional burden of medications in older people. Arch Intern Med 167 (8):781-787.
 442 doi:10.1001/archinte.167.8.781

443 15. Gnjjidic D, Le Couteur DG, Abernethy DR, Hilmer SN (2012) Drug burden index and Beers'
 444 criteria: impact on functional outcomes in older people living in self-care retirement villages.
 445 J Clin Pharmacol 52 (2):258-265. doi:10.1177/0091270010395591

446 16. Duran CE, Azermai M, Vander Stichele RH (2013) Systematic review of anticholinergic
 447 risk scales in older adults. Eur J Clin Pharmacol 69 (7):1485-1496. doi:10.1007/s00228-013-
 448 1499-3

449 17. Boustani M, Campbell N, Munger S, Maidment I, Fox C (2008) Impact of anticholinergics
 450 on the aging brain: a review and practical application. Aging Health 4 (3)

451 18. Ailabouni N, Mangin D, Nishtala PS (2017) Deprescribing anticholinergic and sedative
 452 medicines: protocol for a Feasibility Trial (DEFEAT-polypharmacy) in residential aged care
 453 facilities. BMJ Open 7 (4):e013800. doi:10.1136/bmjopen-2016-013800

454 19. Tune LE (2001) Anticholinergic effects of medication in elderly patients. J Clin Psychiatry
 455 62 Suppl 21:11-14

456 20. Leon AC, Davis LL, Kraemer HC (2011) The role and interpretation of pilot studies in
 457 clinical research. J Psychiatr Res 45 (5):626-629. doi:10.1016/j.jpsychires.2010.10.008

21. Teare MD, Dimairo M, Shephard N, Hayman A, Whitehead A, Walters SJ (2014) Sample size requirements to estimate key design parameters from external pilot randomised controlled trials: a simulation study. *Trials* 15:264. doi:10.1186/1745-6215-15-264
22. Braun V, Clarke V (2006) Using thematic analysis in psychology. *Qualitative Research in Psychology* 3 (2):77-101
23. Cameron RA DT, Richardson S, Ahmed E, Sukumaran A. (2013) Lessons from the field: Applying the Good Reporting of A Mixed Methods Study (GRAMMS) framework'. *Electronic Journal of Business Research Methods* 11 (2):53-64
24. Laycock J, Holmes DM (2003) The place of physiotherapy in the management of pelvic floor dysfunction. *The Obstetrician & Gynaecologist* 5 (4):194-199
25. Anthierens S, Verhoeven V, Schmitz O, Coenen S (2017) Academic detailers' and general practitioners' views and experiences of their academic detailing visits to improve the quality of analgesic use: process evaluation alongside a pragmatic cluster randomized controlled trial. *BMC Health Serv Res* 17 (1):841. doi:10.1186/s12913-017-2797-8
26. Allen M, Ferrier S, O'Connor N, Fleming I (2007) Family physicians' perceptions of academic detailing: a quantitative and qualitative study. *BMC Med Educ* 7:36. doi:10.1186/1472-6920-7-36
27. Soumerai SB, Avorn J (1990) Principles of educational outreach ('academic detailing') to improve clinical decision making. *JAMA* 263 (4):549-556
28. Soumerai SB (1998) Principles and uses of academic detailing to improve the management of psychiatric disorders. *Int J Psychiatry Med* 28 (1):81-96. doi:10.2190/btca-q06p-mgcq-r0l5
29. Thabane L, Hopewell S, Lancaster GA, Bond CM, Coleman CL, Campbell MJ, Eldridge SM (2016) Methods and processes for development of a CONSORT extension for reporting pilot randomized controlled trials. *Pilot Feasibility Stud* 2:25. doi:10.1186/s40814-016-0065-z
30. Bruyndonckx R, Verhoeven V, Anthierens S, Cornelis K, Ackaert K, Gielen B, Coenen S (2018) The implementation of academic detailing and its effectiveness on appropriate prescribing of pain relief medication: a real-world cluster randomized trial in Belgian general practices. *Implement Sci* 13 (1):6. doi:10.1186/s13012-017-0703-8